






## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : <b>C07K 7/16, A61K 37/34</b>		<b>A1</b>	(11) International Publication Number: <b>WO 91/13092</b>
			(43) International Publication Date: <b>5 September 1991 (05.09.91)</b>
(21) International Application Number: <b>PCT/SE91/00154</b>		(74) Agents: <b>NILSSON, Brita et al.; Oscar Grahn Patentbyrå AB, P.O. Box 19540, S-104 32 Stockholm (SE).</b>	
(22) International Filing Date: <b>26 February 1991 (26.02.91)</b>			
(30) Priority data: <b>9000691-7 27 February 1990 (27.02.90) SE</b>		(81) Designated States: <b>AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH (European patent), CM (OAPI patent), DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GA (OAPI patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU (European patent), MC, MG, ML (OAPI patent), MR (OAPI patent), MW, NL (European patent), NO, PL, RO, SD, SE (European patent), SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent), US.</b>	
(71) Applicant (for all designated States except US): <b>FERRING AB [SE/SE]; P.O. Box 30561, S-200 62 Malmö (SE).</b>			
(72) Inventors; and (75) Inventors/Applicants (for US only): <b>AURELL, Carl-Johan [SE/SE]; Erikslustvägen 42 G, S-217 73 Malmö (SE). MELIN, Per [SE/SE]; Svcagatan 100 C, S-216 15 Malmö (SE). NILSSON, Anders [SE/SE]; Korsåkersvägen 37, S-222 50 Lund (SE). TROJNAR, Jerzy [SE/SE]; Stenöregatan 36, S-230 44 Vintrie (SE).</b>		Published With international search report.	

(54) Title: **VASOACTIVE VASOTOCIN DERIVATIVES**

## (57) Abstract

Vasotocin derivatives of formula (I), wherein Hmp = a 2-hydroxy-3-mercaptopropionic acid residue (a), Z = Phe or Tyr, Y = Hgn or Hci, X = (b), Q = H or from 1 to 3 amino acid residues of the same or different natural or unnatural L- or D-amino acids, and n is 1, 2 or 3, are disclosed. A pharmaceutical composition which comprises at least one vasotocin derivative as defined above is intended for use as a vasoconstrictive agent.

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

**FOR THE PURPOSES OF INFORMATION ONLY**

AT	Austria	ES	Spain	MC	Monaco
AU	Australia	FR	France	LU	Luxembourg
BB	Barbados	FI	Finland	LK	Sri Lanka
BE	Belgium	GA	Gabon	LI	Liechtenstein
BF	Burkina Faso	GB	United Kingdom	KR	Republic of Korea
BG	Bulgaria	GN	Guinea	of Korea	
BJ	Benin	GR	Greece	RP	Democratic People's Republic
BR	Brazil	HU	Hungary	JP	Japan
CA	Canada	IT	Italy	SD	Sudan
CF	Central African Republic	RO	Romania	SE	Sweden
CG	Congo	PL	Poland	SN	Senegal
CH	Switzerland	NO	Norway	SU	Soviet Union
CI	Côte d'Ivoire	NL	Netherlands	TC	Togo
CM	Cameroon	MW	Malawi	TD	Chad
CS	Czechoslovakia	MR	Mauritania	US	United States of America
DE	Germany	MN	Mongolia		
DK	Denmark	ML	Mali		

VASOACTIVE VASOTOCIN DERIVATIVES

5

The present invention relates to new vasotocin derivatives, more specifically such vasotocin derivatives as differ from the native hormone in that the vasotocin (VT) structure has been modified at positions 1, 4, 8 and optionally 2.

10

The new VT derivatives are vasoactive, more particularly by specifically raising the blood pressure, and in some cases have a considerably prolonged effect.

15

Background

The peptide hormone vasopressin, produced by the posterior lobe of the pituitary, mainly has two functions, that is the hormone has both an antidiuretic effect (reduced excretion of urine) and a contracting effect on smooth muscles in the vascular wall, the latter effect causing a blood pressure increase and a reduced tendency to bleeding. In clinical use, vasopressin thus has a non-specific effect of short duration.

25

Today, there is on the market a vasopressin analog having a prolonged effect, namely lysine-vasopressin extended in the N-terminal by three amino acid residues. This vasopressin analog acts as a so-called prohormone or hormonogen, i.e. it increases the duration of the vasopressin effect. The

30

extended vasopressin analog has in itself a very small pharmacological effect which does not occur until the extra N-terminal amino acid residues are cleaved by enzymatic hydrolysis and free lysine-vasopressin is formed. Besides the prolonged effect, such a prohormone is advantageous in that

35

the risk of overdosage is minimised by the limited enzyme capacity of the organism determining the plasma levels of the liberated vasopressin. In this manner, it is possible to avoid excessively high plasma levels of vasopressin possibly leading to abnormally increased blood pressure which may harm


the patient. The above-mentioned vasopressin analog however suffers from major drawbacks by having low potency and, like vasopressin, being non-specific.

- 5 There is a need for vasoconstrictive substances for use as bleeding inhibitors and in so-called orthostatic hypotension, i.e. conditions of blood pressure drop following changes of body position. These agents should specifically increase blood pressure, thus having a low antidiuretic effect in  
10 order to avoid water intoxication in patients subjected to long-term treatment. Also, it is advantageous if they exhibit an effect of long duration.

Recently, we have filed (on October 7, 1987) a Swedish patent  
15 application SE 8703855-0 (corresponding to PCT/SE88/00509) comprising vasotocin derivatives having specific blood pressure increasing activity. The vasotocin derivatives according to the present invention differ structurally from the vasotocin derivatives according to said prior Swedish  
20 patent application mainly in that they have a further modification at position 4 of the vasotocin structure, i.e. they have homoglutamine or homocitrulline at position 4.

#### Description of the invention

25 The present new vasoactive vasotocin derivatives specifically increase blood pressure, i.e. they are pressor-specific, meaning a high ratio of blood pressure to antidiuretic activity. In particular the antidiuretic effect (reduced  
30 excretion of urine) of the parent molecule is eliminated. Furthermore they have a considerably prolonged effect in some cases. The compounds according to the invention are intended to be used in a pharmaceutical composition for inhibiting bleeding and in conditions of blood pressure drop following  
35 changes of body position, so-called orthostatic hypotension, and also as general blood pressure increasing agents. The VT derivatives according to the invention are of the formula




5

wherein

Hmp = a 2-hydroxy-3-mercaptopropionic acid residue,


10



Z = phenylalanine (Phe) or tyrosine (Tyr)

Y = homoglutamine (Hgn) or homocitrulline (Hci)

15



20

Q = H or from 1 to 3 amino acid residues of the same or different natural or unnatural L- or D-amino acids, and n is 1, 2 or 3.

25

The VT derivatives according to the invention can be presented in the form of pharmaceutical compositions in which at least one VT derivative according to the invention is included as active ingredient, together with pharmaceutically acceptable additives and/or diluents. The pharmaceutical compositions according to the invention preferably are in the form of preparations suitable for parenteral administration. They are suitably administered by injection, infusion or intranasal application. The diluent may be e.g. a physiological saline solution.

30

35

A pharmaceutical composition according to the invention may contain a specifically blood pressure increasing derivative having a relatively short duration for providing an instant effect, in combination with a specifically blood pressure

increasing derivative having a long duration for providing a prolongation of the effect.

Preparation of the VT derivatives according to the invention

5

The VT derivatives according to the invention can be prepared by methods analogous with those which are known in the peptide field.

- 10 For instance, the compounds according to the invention can be prepared in conventional manner by coupling amino acids stepwise to one another in liquid phase, e.g. as disclosed by Law, H.B. & Du Vigneaud, V. in Journal of the American Chemical Society 82, (1960) 4579-4581, Zhuze, A.L., Jöst, K.,
- 15 Kasafi'rek, E. & Rudinger, J. in Collection of Czechoslovak Chemical Communications 29 (1964), 2648-2662, and modified by Larsson, L.-E., 5 Lindeberg, G., Melin, P. / Pliška, V. in Journal of Medicinal Chemistry 21, (1978), 352-356.
- 20 The coupling of the amino acids to one another, yielding so-called peptide bonds, can also be effected with a solid phase (generally a resin) as starting material to which the C-terminal of the first amino acid is coupled, whereupon the C-terminal of the next amino acid is coupled to the N-terminal of the first amino acid and so on. Finally, the
- 25 finished peptide is liberated from the solid phase. In the Examples hereinbelow, this so-called solid phase technique has been used in accordance with the disclosure of Merrifield, R.B., J. Am. Chem. Soc. (1963) 85, 2149, Merrifield, R.B. Biochemistry (1964), 3, 1385 and König, W., Geiger, R.,
- 30 Chem. Ber. (1970), 103, 788.

General description of synthesis

- 35 All the VT derivatives prepared in the Examples given below were synthesised on an Applied Biosystems 430A Peptide Synthesizer using a double coupling program with a termination step after the second coupling. The resin used was of 4-methylbenzhydrylamine type with a theoretical loading

- of 0.65 meq/g (Peninsula Laboratories Inc., USA). The final product of the synthesis was dried in vacuo overnight. The peptide was then cleaved from the resin by treatment with liquid hydrogen fluoride in the presence of anisole and ethyl-methyl-sulphide as scavengers (HF:anisole:EMS - 10:05:05). After removal of hydrogen fluoride by evaporation, the resin was suspended in ethyl acetate (100 ml) and filtered. The solid was washed on filter with additional ethyl acetate (3x100 ml), and the cleaved peptide was extracted with acetic acid (100 ml). The extract was promptly diluted to a volume of 1.5 l with 20% acetic acid in methanol and treated with 0.1 M solution of iodine in methanol until a faint brown colour remained. Then a Dowex 1x8 ion exchanger in acetate form (15 g) (Bio-Rad, Richmond, CA) was added and the mixture filtered. The filtrate was evaporated and the residue freeze-dried from water. The product was then purified by reversed phase liquid chromatography on a column filled with Kromasil® 13 µ (EKA Nobel, Surte, Sweden) in a suitable system containing acetonitrile in 0.1% trifluoroacetic acid water solution. The samples collected from the column were analysed by analytical high performance liquid chromatography (HPLC) (Spectra Physics Inc. USA 8800) equipped with a Vydac 5 µ C18 column (Vydac Inc., USA). Fractions containing pure substance were pooled, the solvent was evaporated and the product freeze-dried from water. The final HPLC analysis was performed on ready product, and the structure of the peptide was confirmed by amino acid analysis and fast atom bombardment mass spectrometry (FAB MS).
- All amino acids used during the synthesis were L-amino acids, and they were protected with a tert-butoxy-carbonyl group at the α-amino function. The side chains were protected as follows:
- Hmp(Mob), Cys(Mob), Dab(Cbz).

The abbreviations within brackets are:

- Cbz = carbobenzoxy;  
Mob = 4-methoxybensyl; and  
5 Boc = t-butyloxycarbonyl

The amino acid derivatives were supplied by Bachem AG, Switzerland.

10 Further abbreviations used are:

- Dab = L-2,4-diaminobutyric acid  
Abu = L-2-aminobutyric acid  
Hgn = homoglutamine  
15 Hci = homocitrulline  
Hmp = 2-hydroxy-3-mercaptopropionic acid  
OPfp = pentafluorophenyl ester  
DIPEA = diisopropylethylamine

20 EXAMPLE 1

1-Hmp-2-Phe-4-Hgn-8-Dab-VT  
[n = 2 and Q = H ]

25 The peptide was synthesised according to the general description. 2-hydroxy-mercaptopropionic acid[S-(p-methoxy)benzyl] was used for position 1. Purity (HPLC): 99.5% (18.4% acetonitrile in 0.1% TFA, retention time 9.13 min at 1.5 ml/min, detection at 223 nm).

30 The structure was confirmed by amino acid analysis and FAB MS analysis.

EXAMPLE 2

1-Hmp-2-Phe-4-Hgn-8-Dab(Ala)-VT  
35 [n = 2 and Q = Ala]

The oxidized and purified nonapeptide Hmp-Phe-Ile-Hgn-Asn-Cys-Pro-Dab-Gly-NH<sub>2</sub> (150 mg; prepared by solid phase method

according to the general description) was dissolved in DMF (2 ml) and previously formed Boc-Ala-OPfp (4 equivalents) was added and pH was adjusted to 8-8.5 (DIPEA). The reaction mixture was stirred overnight at room temperature. The product was isolated by precipitation with ethyl acetate, filtration and drying in vacuo.

The product was then treated with TFA/CH<sub>2</sub>Cl<sub>2</sub> 1:1 (20 ml), stirred for 30 min, evaporated and then treated with diethyl ether (100 ml). The precipitation was separated by filtration and dried in vacuo.

The product was purified by reversed phase liquid chromatography.

Purity (HPLC): 99.8% (17.6% acetonitrile in 0.1% TFA, retention time 8.56 min at 2 ml/min, detection at 223 nm).

The structure was confirmed by amino acid analysis and FAB MS analysis.

### EXAMPLE 3

1-Hmp-2-Phe-4-Hgn-8-Dab(Abu)-VT

[n = 2 and Q = Abu]

The oxidized and purified nonapeptide Hmp-Phe-Ile-Hgn-Asn-Cys-Pro-Dab-Gly-NH<sub>2</sub> (100 mg; prepared by solid phase method according to the general description) was dissolved in DMF (2 ml) and previously formed Boc-Abu-OPfp (4 equivalents) was added and pH was adjusted to 8-8.5 (DIPEA). The reaction mixture was stirred overnight at room temperature. The product was isolated by precipitation with ethyl acetate, filtration and drying in vacuo.

The product was then treated with TFA/CH<sub>2</sub>Cl<sub>2</sub> 1:1 (20 ml), stirred for 30 min, evaporated and then treated with diethyl ether (100 ml). The precipitation was separated by filtration and dried in vacuo.

The product was purified by reversed phase liquid chromatography.

- 5 Purity (HPLC): 99.5% (17.6% acetonitrile in 0.1% TFA, retention time 9.82 min at 2 ml/min, detection at 223 nm).

The structure was confirmed by amino acid analysis and FAB MS analysis.

10

EXAMPLE 4

1-Hmp-2-Phe-4-Hci-8-Dab-VT

[n = 2 and Q = H ]

- 15 The peptide was synthesised according to the general description. 2-hydroxy-mercaptopropionic acid[S-(p-methoxy)benzyl] was used for position 1. Purity (HPLC): 98.5% (17.6% acetonitrile in 0.1% TFA, retention time 10.68 min at 2 ml/min, detection at 223 nm).

20

The structure was confirmed by amino acid analysis and FAB MS analysis.

EXAMPLE 5

- 25 1-Hmp-2-Phe-4-Hci-8-Dab(Abu)-VT

[n = 2 and Q = Abu]

- 30 The oxidized and purified nonapeptide Hmp-Phe-Ile-Hci-Asn-Cys-Pro-Dab-Gly-NH<sub>2</sub> (100 mg; prepared by solid phase method according to the general description) was dissolved in DMF (2 ml) and previously formed Boc-Abu-OPfp (4 equivalents) was added and pH was adjusted to 8-8.5 (DIPEA). The reaction mixture was stirred overnight at room temperature. The product was isolated by precipitation with ethyl acetate, filtration and drying in vacuo.
- 35

The product was then treated with TFA/CH<sub>2</sub>Cl<sub>2</sub> 1:1 (20 ml), stirred for 30 min, evaporated and then treated with diethyl

ether (100 ml). The precipitation was separated by filtration and dried in vacuo.

5 The product was purified by reversed phase liquid chromatography.

Purity (HPLC): 99.5% (20.0% acetonitrile in 0.1% TFA, retention time 5.94 min at 2 ml/min, detection at 223 nm).

10 The structure was confirmed by amino acid analysis and FAB MS analysis.

#### EXAMPLE 6

1-Hmp-4-Hgn-8-Orn-VT

15 [n = 2 and Q = H ]

The peptide was synthesised according to the general description. 2-hydroxy-mercaptopropionic acid[S-(p-methoxy)benzyl] was used for position 1. Purity (HPLC): 98.5% (14.4% acetonitrile in 0.1% TFA, retention time 5.83 min at 2 ml/min, detection at 223 nm).

20

The structure was confirmed by amino acid analysis and FAB MS analysis.

25

#### EXAMPLE 7

1-Hmp-4-Hgn-8-Dab-VT

[n = 2 and Q = H ]

30 The peptide was synthesised according to the general description. 2-hydroxy-mercaptopropionic acid[S-(p-methoxy)benzyl] was used for position 1. Purity (HPLC): >99% (16.0% acetonitrile in 0.1% TFA, retention time 4.98 min at 2 ml/min, detection at 223 nm).

35

The structure was confirmed by amino acid analysis and FAB MS analysis.

Pharmacological tests

5 Vasotocin derivatives according to the invention have been tested for potency of both blood pressure and antidiuretic activity in a so-called 4-point test, i.e. the activity of the test substances has been related to a standard preparation (AVP = argininevasopressin), and the effects of two dose levels for each substance have been analysed. In addition, three pressor-specific VT derivatives according to our previous application SE 8703855-0 have been tested for a comparison, namely 1-Hmp-2-Phe-8-Orn-VT (compound 2 in Table 1), 1-Hmp-2-Phe-8-Dab-VT (compound 3 in Table 1), and 1-Hmp-2-Phe-8-Dab(Ala)-VT (compound 5 in Table 1).

15 Blood pressure tests were carried out on anaesthetised Sprague Dawley rats (about 250 g), previously treated with dibenamine (Dekanski, J., 1952. Br. J. Pharmacol. 7, 567). Maximal blood pressure increase after intravenous injections of peptide was used as a measure of the effect, expressed as intensity.

In addition to potency determination based on effect intensity, a measure of the length of the effect has been stated (index of persistence (I.P.), Pliška, V., 1966. 25 *Arzheim. Forsch.* 16, 886). This dimensionless factor is a measure of the effect duration of the respective analog in relation to the standard AVP.

30 Antidiuretic potency was determined with the aid of anaesthetised water-loaded Sprague Dawley rats (200 g) (Larsson, L.E., Lindeberg, G., Melin, P. and Pliška, V., 1978, *J. Med. Chem.* 21, 353). Maximal increase of urine conductivity after intravenous injections was used as effect parameter.

In these two tests, a comparison was made between the effects of the respective derivative and the effect of a standard

preparation, AVP, and potency was determined with the aid of a 4-point test and is indicated in international units per micromole (IU/ $\mu$ mole) (Stürmer, E., in Handbook of Experimental Pharmacology, 1966, Vol 23, pp 130-189).

5

The specificity in respect of blood pressure is indicated by the ratio of potency blood pressure/potency antidiuresis (BP/AD).

10 The pharmacological results are given in Table 1.

From Table 1 it appears that the compounds according to the invention retain a very high potency in respect of blood pressure increase and effect duration.

15

By the introduction of homoglutamine or homocitrulline at position 4 the antidiuretic activity has been practically eliminated. Thus, the present invention is unique in that the pressor specificity (ratio of blood pressure to antidiuretic activity) has been increased approximately 2 to 10 times in comparison to the already pressor specific derivatives of our SE 8703855-0 (modifications at positions 1, 2 and 8 of the parent molecule; see Table 1).

20

25 The combination of this modification with previously made substitutions at positions 1, 2 and 8 has led to analogs with high potency, long duration of action and extreme pressor specificity. Thus, based on the animal experiments presented, the new substances may, in therapy be expected to completely lack any water accumulating effect (antidiuretic), thus  
30 totally avoiding the risk of water intoxications of the patients.

35

Example of the preparation of a pharmaceutical composition

35

The VT derivative is dissolved in distilled water together with mannitol. The solution is poured into an ampoule,

subjected to freeze-drying and sealed. The contents in the ampoule can then when desired, be diluted with an isotonic saline solution to a concentration suitable for administration.


TABLE 1

ANALOG	BLOOD PRESSURE BP		ANTIDIURESIS AD IU/ $\mu$ mole	BP/AD (measure of specificity)
	IU/ $\mu$ mole	I.P (measure of duration of effect)		
1. AVP (Reference)	614 $\pm$ 25	1.0	620 $\pm$ 54	1.0
2. 1-Hmp-2-Phe-8-Orn-VT	421 $\pm$ 41	3.1 $\pm$ 1.3	9.4 $\pm$ 1.1	45
3. 1-Hmp-2-Phe-8-Dab-VT	657 $\pm$ 32	2.7 $\pm$ 0.6	18 $\pm$ 2.1	37
4. 1-Hmp-2-Phe-4-Hg -8-Dab-VT (Ex. 1)	214 $\pm$ 6	2.3 $\pm$ 0.5	0.3 $\pm$ 0.02	713
5. 1-Hmp-2-Phe-8-Dab(Ala)-VT	360 $\pm$ 31	6.7 $\pm$ 1.5	4.3 $\pm$ 0.3	84
6. 1-Hmp-2-Phe-4-Hg -8-Dab(Ala)-VT (Ex. 2)	117 $\pm$ 6	6.6 $\pm$ 1.0	0.2 $\pm$ 0.02	585
7. 1-Hmp-2-Phe-4-Hgn-8-Dab(Abu)-VT (Ex. 3)	163 $\pm$ 15	8.6 $\pm$ 2.3	0.2 $\pm$ 0.01	858
8. 1-Hmp-2-Phe-4-Hci-8-Dab-VT (Ex. 4)	157 $\pm$ 13	1.7 $\pm$ 0.05	0.3 $\pm$ 0.05	523
9. 1-Hmp-2-Phe-4-Hci-8-Dab(Abu)-VT (Ex. 5)	67 $\pm$ 5	3.6 $\pm$ 0.7	0.2 $\pm$ 0.04	335
10. 1-Hmp-4-Hgn-8-Orn-VT (Ex. 6)	473 $\pm$ 62	7.3 $\pm$ 1.8	1.0 $\pm$ 0.2	473
11. 1-Hmp-4-Hgn-8-Dab-VT (Ex. 7)	536 $\pm$ 53	3.8 $\pm$ 0.8	2.9 $\pm$ 0.5	185

CLAIMS

5


1. Vasotocin derivative of the formula



wherein

Hmp = a 2-hydroxy-3-mercaptopropionic acid residue

15




Z = phenylalanine (Phe) or tyrosine (Tyr)

20

Y = homoglutamine (Hgn) or homocitrulline (Hci)

X = -HN-CH-CO-



25

Q = H or from 1 to 3 amino acid residues of the same or  
different natural or unnatural L- or D-amino acids, and  
n is 1, 2 or 3.

30

2. Vasotocin derivative as claimed in claim 1,  
characterised in that

Z = Phe,

Y = Hgn,

n is 2, and

35

Q is H.

3. Vasotocin derivative as claimed in claim 1,  
c h a r a c t e r i s e d    i n t h a t

5    Z = Phe,  
     Y = Hgn,  
     n is 2, and  
     Q is alanyl.

10    4. Vasotocin derivative as claimed in claim 1,  
     c h a r a c t e r i s e d    i n t h a t

     Z = Phe,  
     Y = Hgn,  
     n is 2, and  
15    Q is L-2-aminobutyryl.

5. Vasotocin derivative as claimed in claim 1,  
c h a r a c t e r i s e d    i n t h a t

     Z = Phe,  
20    Y = Hci,  
     n = 2, and  
     Q = H.

6. Vasotocin derivative as claimed in claim 1,  
25    c h a r a c t e r i s e d    i n t h a t

     Z = Tyr,  
     Y = Hgn,  
     n = 3, and  
     Q = H.

30

7. Vasotocin derivative as claimed in claim 1,  
c h a r a c t e r i s e d    i n t h a t

     Z = Tyr,  
     Y = Hgn,  
35    n = 2, and  
     Q = H.

8. Pharmaceutical composition, c h a r a c t e r i s e d  
in that it comprises at least one vasotocin derivative as  
claimed in claim 1 as active ingredient, together with  
5 pharmaceutically acceptable additive(s) and/or diluent(s).




9. Pharmaceutical composition as claimed in claim 8,  
c h a r a c t e r i s e d in that it is in the form  
of a preparation suitable for parenteral administration.  
10

10. Pharmaceutical composition as claimed in claim 9,  
c h a r a c t e r i s e d in that it is in the form  
of a solution suitable for intranasal administration.

15

# INTERNATIONAL SEARCH REPORT

International Application No PCT/SE 91/00154

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (If several classification symbols apply, indicate all) <sup>6</sup> According to International Patent Classification (IPC) or to both National Classification and IPC <b>IPC5: C 07 K 7/16, A 61 K 37/34</b>														
<b>II. FIELDS SEARCHED</b> <div style="text-align: right; margin-right: 100px;">Minimum Documentation Searched<sup>7</sup></div> <table style="width: 100%; border: none;"> <tr> <td style="width: 20%; border: none;">Classification System</td> <td style="border: none;">Classification Symbols</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px;">IPC5</td> <td style="border: 1px solid black; padding: 5px;">A 61 K; C 07 K</td> </tr> </table> <div style="text-align: center; margin-top: 5px;">Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in Fields Searched<sup>8</sup></div>			Classification System	Classification Symbols	IPC5	A 61 K; C 07 K								
Classification System	Classification Symbols													
IPC5	A 61 K; C 07 K													
SE,DK,FI,NO classes as above														
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup></b> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 10%;">Category<sup>*</sup></th> <th style="width: 60%;">Citation of Document,<sup>11</sup> with indication, where appropriate, of the relevant passages<sup>12</sup></th> <th style="width: 30%;">Relevant to Claim No.<sup>13</sup></th> </tr> </thead> <tbody> <tr> <td style="text-align: center; vertical-align: top;">A</td> <td>WO, A1, 8903393 (FERRING AB) 20 April 1989, see the whole document --</td> <td style="text-align: center; vertical-align: top;">1-10</td> </tr> <tr> <td style="text-align: center; vertical-align: top;">A</td> <td>Br. J. Pharmac., Vol. 67, 1979 G.W. Bisset et al.: "Hydroxy Analogues of oxytocin and of lysinevasopressin", see page 575 - page 585 --</td> <td style="text-align: center; vertical-align: top;">1-10</td> </tr> <tr> <td style="text-align: center; vertical-align: top;">A</td> <td>Endocrinology, Vol. 112, No. 1, 1983 Dean W. Cheesman et al.: "Anovulatory Effect of Synthetic Analogs of Arginine Vasotocin in the Rat", see page 269 - page 276 -- -----</td> <td style="text-align: center; vertical-align: top;">1-10</td> </tr> </tbody> </table>			Category <sup>*</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>	A	WO, A1, 8903393 (FERRING AB) 20 April 1989, see the whole document --	1-10	A	Br. J. Pharmac., Vol. 67, 1979 G.W. Bisset et al.: "Hydroxy Analogues of oxytocin and of lysinevasopressin", see page 575 - page 585 --	1-10	A	Endocrinology, Vol. 112, No. 1, 1983 Dean W. Cheesman et al.: "Anovulatory Effect of Synthetic Analogs of Arginine Vasotocin in the Rat", see page 269 - page 276 -- -----	1-10
Category <sup>*</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>												
A	WO, A1, 8903393 (FERRING AB) 20 April 1989, see the whole document --	1-10												
A	Br. J. Pharmac., Vol. 67, 1979 G.W. Bisset et al.: "Hydroxy Analogues of oxytocin and of lysinevasopressin", see page 575 - page 585 --	1-10												
A	Endocrinology, Vol. 112, No. 1, 1983 Dean W. Cheesman et al.: "Anovulatory Effect of Synthetic Analogs of Arginine Vasotocin in the Rat", see page 269 - page 276 -- -----	1-10												
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p><sup>*</sup> Special categories of cited documents: <sup>10</sup></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&amp;" document member of the same patent family</p> </div> </div>														
<b>IV. CERTIFICATION</b> <table style="width: 100%; border: none;"> <tr> <td style="width: 50%; border: none;">Date of the Actual Completion of the International Search</td> <td style="width: 50%; border: none;">Date of Mailing of this International Search Report</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px;">27th May 1991</td> <td style="border: 1px solid black; padding: 5px;">1991 -05- 05</td> </tr> <tr> <td style="border: none;">International Searching Authority</td> <td style="border: none;">Signature of Authorized Officer</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px; text-align: center;">SWEDISH PATENT OFFICE</td> <td style="border: 1px solid black; padding: 5px;">             Elisabeth Carlborg         </td> </tr> </table>			Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	27th May 1991	1991 -05- 05	International Searching Authority	Signature of Authorized Officer	SWEDISH PATENT OFFICE	 Elisabeth Carlborg				
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report													
27th May 1991	1991 -05- 05													
International Searching Authority	Signature of Authorized Officer													
SWEDISH PATENT OFFICE	 Elisabeth Carlborg													

ANNEX TO THE INTERNATIONAL SEARCH REPORT  
ON INTERNATIONAL PATENT APPLICATION NO.PCT/SE 91/00154

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.  
The members are as contained in the Swedish Patent Office EDP file on 91-04-30  
The Swedish Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A1- 8903393	89-04-20	AU-D- 2533988 EP-A- 0380554 SE-A- 8703855	89-05-02 90-08-08 89-04-08